

Effect of Number of Sprints in an SIT Session on Change in $\dot{V}O_{2\max}$: A Meta-analysis

NIELS B. J. VOLLAARD¹, RICHARD S. METCALFE², and SEAN WILLIAMS¹

¹Department for Health, University of Bath, Bath, UNITED KINGDOM; and ²School of Sport, Ulster University, Northern Ireland, UNITED KINGDOM

ABSTRACT

VOLLAARD, N. B. J., R. S. METCALFE, and S. WILLIAMS. Effect of Number of Sprints in an SIT Session on Change in $\dot{V}O_{2\max}$: A Meta-analysis. *Med. Sci. Sports Exerc.*, Vol. 49, No. 6, pp. 1147–1156, 2017. **Purpose:** Recent meta-analyses indicate that sprint interval training (SIT) improves cardiorespiratory fitness ($\dot{V}O_{2\max}$), but the effects of various training parameters on the magnitude of the improvement remain unknown. The present meta-analysis examined the modifying effect of the number of sprint repetitions in an SIT session on improvements in $\dot{V}O_{2\max}$. **Methods:** The databases PubMed and Web of Science were searched for original studies that have examined pre- and posttraining $\dot{V}O_{2\max}$ in adults after ≥ 2 wk of training consisting of repeated (≥ 2) Wingate-type cycle sprints, published up to May 1, 2016. Articles were excluded if they were not in English; if they involved patients, athletes, or participants with a mean baseline $\dot{V}O_{2\max}$ of >55 mL \cdot kg⁻¹ \cdot min⁻¹ or a mean age <18 yr; and if an SIT trial was combined with another intervention or used intervals shorter than 10 s. A total of 38 SIT trials from 34 studies were included in the meta-analysis. Probabilistic magnitude-based inferences were made to interpret the outcome of the analysis. **Results:** The meta-analysis revealed a likely large effect of a typical SIT intervention on $\dot{V}O_{2\max}$ (mean \pm 90% confidence limits = $7.8\% \pm 4.0\%$) with a possibly small modifying effect of the maximum number of sprint repetitions in a training session ($-1.2\% \pm 0.8\%$ decrease per two additional sprint repetitions). Apart from possibly small effects of baseline $\dot{V}O_{2\max}$ and age, all other modifying effects were unclear or trivial. **Conclusion:** We conclude that the improvement in $\dot{V}O_{2\max}$ with SIT is not attenuated with fewer sprint repetitions, and possibly even enhanced. This means that SIT protocols can be made more time efficient, which may help SIT to be developed into a viable strategy to impact public health. **Key Words:** SYSTEMATIC REVIEW, CARDIORESPIRATORY FITNESS, AEROBIC CAPACITY, SPRINT INTERVAL TRAINING

The global increase in prevalence of noncommunicable diseases for the past decades (34) can be attributed, at least in part, to the low levels of physical activity undertaken by the majority of the general population (16). In light of this, a key aim of public health organizations is to increase population physical activity levels (20). Of the health markers that can be improved by physical activity, maximal aerobic capacity ($\dot{V}O_{2\max}$) is consistently shown to be the strongest prognostic marker for future cardiovascular health and premature death in cross-sectional studies (38,56). Furthermore, longitudinal studies demonstrate that improvements in $\dot{V}O_{2\max}$ are associated with substantial reductions in all-cause and cardiovascular mortality during follow-up (9,43).

For the past two decades, relatively high volumes of moderate-intensity aerobic exercise (total time commitment

≥ 150 min \cdot wk⁻¹) have consistently been recommended for improving health markers (20). However, uptake of and adherence to these recommendations remains low in the general population (25), with lack of time identified as one of the main perceived barriers to becoming and remaining physically active (39,41,71). Therefore, the seminal finding by Burgomaster et al. (12) that a training protocol consisting of repeated brief all-out cycle sprints (i.e., Wingate sprints) is associated with aerobic adaptations has led to substantial interest in the use of (sub)maximal high-intensity interval training (HIIT) and supramaximal sprint interval training (SIT) as time-efficient alternative/adjunct exercise strategies for improving $\dot{V}O_{2\max}$ (21). The most commonly studied SIT protocol consists of four to seven repeated 30-s Wingate sprints, thus resulting in less than 4 min of high-intensity exercise per session (75). For the past few years, several meta-analyses have reported the efficacy of SIT in increasing $\dot{V}O_{2\max}$ (24,53,65,75). These have concluded that in healthy individuals, SIT improves $\dot{V}O_{2\max}$ to a similar (24) or greater extent (53) than traditional aerobic training, with greater benefits for individuals with lower pretraining $\dot{V}O_{2\max}$ (53,75).

Although these findings provide strong support for the effectiveness of SIT in improving $\dot{V}O_{2\max}$, surprisingly few efforts have been made to identify “optimal” SIT protocols, e.g., protocols which will either provide the greatest increase

Address for correspondence: Niels B.J. Vollaard, Ph.D., Department for Health, University of Bath, Bath, BA2 7AY, United Kingdom; E-mail: n.vollaard@stir.ac.uk.

Submitted for publication October 2016.

Accepted for publication January 2017.

0195-9131/17/4906-1147/0

MEDICINE & SCIENCE IN SPORTS & EXERCISE®

Copyright © 2017 by the American College of Sports Medicine

DOI: 10.1249/MSS.0000000000001204

in $\dot{V}O_{2\max}$, or a set increase with the lowest total training volume or time commitment. Weston et al. (75) reported a likely small effect of increasing the intervention duration and a possibly moderate effect of increasing the work-to-rest ratio, but no studies have meta-analyzed or directly investigated the potential effects of the number of sprint repetitions in an SIT session. This parameter is particularly important as it has a large influence on the total duration of a training session, as well as the level of fatigue (44) and affective responses (19) experienced by the participant, thus influencing the likelihood of individuals taking up and adhering to a specific SIT intervention (26). As the main aim of investigating SIT protocols is generally to identify a time-efficient alternative to aerobic exercise, there is a need to identify the effect of this training parameter on the associated increase in $\dot{V}O_{2\max}$. Recent evidence suggests that the positive effects of SIT on $\dot{V}O_{2\max}$ can be attained with fewer sprints (22,23,35,50), and therefore the aim of the present study was to perform a meta-analysis to provide estimates of the modifying effect of the number of sprint repetitions in SIT protocols on the increase in $\dot{V}O_{2\max}$ in untrained adult participants after training.

METHODS

Literature search criteria and study selection. This study was undertaken in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement guidelines (54). We aimed to identify all studies that have examined pre- and posttraining $\dot{V}O_{2\max}$ after a period of at least 2 wk of training consisting of repeated (≥ 2) all-out Wingate cycle sprints or modifications thereof (e.g., studies using 10-, 15-, or 20-s all-out sprints instead of 30-s Wingate sprints). For this purpose, the electronic databases PubMed and Web of Science were searched for relevant available records up to May 1, 2016, using the 28 possible combinations of the independent variable search terms “Wingate,” “all-out,” “sprint,” and “interval training,” and the dependent variable search terms “fitness,” “aerobic capacity,” “aerobic power,” “ $\dot{V}O_{2\max}$,” “ $\dot{V}O_{2\text{peak}}$,” “oxygen uptake,” and “oxygen consumption.” Relevant studies cited in recent meta-analyses were also used (24,53,65,75), as well as our own recent work (52). The following articles were excluded: 1) review articles/commentaries; 2) articles not written in English; 3) studies concerning patients, athletes, or participants with a mean baseline $\dot{V}O_{2\max}$ of $>55 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ or a mean age <18 yr; 4) animal studies; 5) study trials in which SIT was combined with another intervention; and 6) SIT studies using noncycling exercise, intervals shorter than 10 s, or intervals that were not “all-out.” Two authors (NBJV and RSM) independently conducted the literature search and data extraction, and any discrepancies were resolved by consensus. The reviewers were not blinded to manuscript journals or authors. After removal of duplicate records, the titles and abstracts of all identified articles were screened for records that were

clearly not relevant. These articles were omitted before assessing the full-text versions of the remaining articles for eligibility to be included in the meta-analysis. If more than one article reported data for the same experiment, duplicate data for these participants were only included once. The final data set included the results of 38 trials from 34 studies (Fig. 1).

Data extraction. Full articles were assessed for mean absolute pre- and posttraining $\dot{V}O_{2\max}$. Absolute $\dot{V}O_{2\max}$ ($\text{L}\cdot\text{min}^{-1}$) was used rather than relative $\dot{V}O_{2\max}$ ($\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) as this provides an estimate of true changes in the ability to take up and use oxygen, independent of possible concomitant changes in body mass. Relative $\dot{V}O_{2\max}$ was used for the five studies for which absolute $\dot{V}O_{2\max}$ data were not available (8,42,48,57,68). Any data for $\dot{V}O_{2\max}$ obtained at intermediate time points during the intervention were excluded. The corresponding authors of articles without the required data were contacted by e-mail; authors from 23 studies were contacted (1,2,5,6,10–13,22,23,27,28,31,32,35,40,49,57,62–64,68,70,77), and we received raw data from 17 studies (5,10–13,22,23,27,28,31,40,49,57,62–64,77). Graph digitizer software (DigitizeIt, Braunschweig, Germany) was used to obtain the data from one study for which absolute pre- and posttraining $\dot{V}O_{2\max}$ data were only available in a figure (70). For trials with a no-exercise control group, the effect entered into the meta-analysis was intervention minus control. Data for aerobic exercise comparator groups were not included in the meta-analysis. The effect of training was expressed as a percentage change score. Percentage effects of SIT on $\dot{V}O_{2\max}$ were converted to factors ($= 1 + \text{effect}/100$), log transformed for the analysis, and then back transformed to percentages. Effects were weighted using percentage standard errors derived from exact *P* values, or from estimated errors of measurement as recommended by Weston et al. (75). Under the assumption that studies with similar test protocols and subject characteristics would have similar typical errors of measurement, the typical errors from these studies were averaged (via the weighted mean variance) and assigned to the studies that did not report an exact *P* value (1,2,6,35,46,68,70). The SE was then calculated via the relationship between typical error and SE (72). Finally, data for the following potential moderators were extracted for each study: participant characteristics (sex, age, body mass index, and baseline $\dot{V}O_{2\max}$), training parameters (intervention duration, total number of training sessions, maximal number of sprint repetitions per training session, sprint duration, sprint/recovery ratio, and sprint resistance), and study type (controlled/uncontrolled; dummy variable).

Statistical analysis. To evaluate the extent of publication bias, a funnel plot of model residuals versus their corresponding standard errors was inspected for evidence of asymmetrical scatter (75). This approach takes into account any heterogeneity explained by the meta-regression, which is not accounted for in standard funnel plots of observed effects versus their standard errors. No evidence of asymmetrical scatter was apparent (Fig. 2).




FIGURE 1—Flow diagram of the study selection process.

A mixed effects meta-regression model was conducted using the “metafor” package in R (version 3.2.4, R Foundation for Statistical Computing, Vienna, Austria) (73). The overall effect of SIT on $\dot{V}O_{2max}$ was evaluated using the mean values of the covariates. The modifying effects of covariates were evaluated as the difference between levels (e.g., male/female) for nominal variables. For numeric variables, effects were evaluated as the change in $\dot{V}O_{2max}$ associated with a 2 SD change in the predictor (i.e., a typically low vs a typically high value [33]), or a practically relevant value (e.g., three additional SIT sessions would typically constitute an additional week of training). The random effects in the model specified a between-study SD, representing the typical difference in the true value of the effect in different study settings, plus a within-study random effect to account for within-study repeated measurements (a control treatment and/or more than one training treatment) (75). The SD was doubled before interpreting its magnitude with the scale used to interpret fixed effects (66), for the same reason that the magnitude of the effect of a linear covariate is evaluated with two SD of the covariate (33). We performed a sensitivity analysis to determine whether the inference relating to the modifying effect of maximum number of sprints was substantially altered when two potentially influential studies (with 12 and 15 maximum sprints, respectively [32,64]) were removed from the analysis.

We used magnitude-based inferences to provide an interpretation of the real-world relevance of the outcomes. Uncertainty in effect estimates was expressed as $\pm 90\%$ confidence limits, and as the likelihood that the true value

was beneficial, trivial, or harmful in relation to threshold values for benefit (improved fitness) and harm (reduced fitness) (32). The overall effect of SIT on $\dot{V}O_{2max}$ was interpreted as a clinical outcome, whereby an effect was deemed unclear if the chance that the true value was beneficial was $>25\%$, with odds of benefit relative to odds of harm (odds ratio) of <66 . Modifying effects were evaluated mechanistically and deemed unclear if the likelihood that the




FIGURE 2—Funnel plot of model residuals versus their corresponding standard errors, with 90% confidence interval region.

true value was beneficial and harmful was both >5%. Otherwise, the effect was deemed clear and was qualified with a probabilistic term using the following scale: <0.5%, most unlikely; 0.5%–5%, very unlikely; 5%–25%, unlikely; 25%–75%, possible; 75%–95%, likely; 95%–99.5%, very likely; >99.5%, most likely. As robust anchors for the smallest worthwhile clinical and practical effects relating to $\dot{V}O_{2\max}$ were not available, standardized effect thresholds of 0.2, 0.6, and 1.2 SD were adopted for small, moderate, and large effects, respectively (75). Here, the SD related to the average between-subject variances for baseline $\dot{V}O_{2\max}$; these corresponded to magnitude thresholds of 1.0%, 2.9%, and 5.8%.

RESULTS

Data available for the 34 studies and 38 trials included in the meta-analysis are shown in Table 1 and Figure 3. The meta-analysis indicated an overall likely large effect of an “average” SIT protocol on $\dot{V}O_{2\max}$ (mean \pm 90% confidence limits percent increase in $\dot{V}O_{2\max}$ = 7.8% \pm 4.0%; Table 2). A possibly small effect was evident for the modifying effect of the maximum number of sprint repetitions in a training session ($-1.2\% \pm 0.8\%$ decrease per two additional sprint repetitions; Fig. 4A). The percentage chances that the modifying effect was negative, trivial, or positive were calculated to be 62.7%, 37.3%, and 0.0%, respectively. There were possibly small effects of baseline $\dot{V}O_{2\max}$ ($-1.5\% \pm 1.9\%$ decrease per 10 mL·kg⁻¹·min⁻¹ higher baseline $\dot{V}O_{2\max}$; Fig. 4B) and age ($-1.1\% \pm 1.2\%$ decrease per 7 yr increase; Fig. 4C). All other modifying effects (intervention duration, number of sessions, sprint duration, recovery time, sprint resistance, body mass index, sex, and study type) were unclear or trivial (Table 2). Unexplained variance between studies was 2.2% \pm 0.8% (likely moderate). The inference relating to the effect of maximum number of sprint repetitions was not altered when the two studies with the highest number of sprint repetitions (32,64) were removed from the analysis ($-1.0\% \pm 1.1\%$; possibly small decrease; chances that the modifying effect was negative, trivial, or positive of 51.6%, 48.2%, and 0.0%, respectively).

DISCUSSION

The main aim of the present meta-analysis was to examine the modifying effect of the number of sprint repetitions in an SIT session on the increase in $\dot{V}O_{2\max}$ after training. Using data from 34 training studies and 418 participants, we demonstrate that the improvement in $\dot{V}O_{2\max}$ with SIT is not attenuated with fewer sprint repetitions, and possibly even enhanced. Considering the low physical activity levels in the general population (25), and the fact that lack of time is consistently identified as one of the main perceived barriers to becoming and remaining physically active (39,41,71), this finding has implications for the design of practical SIT

interventions for improving general health. SIT protocols have the potential to be the most time-efficient interventions that are associated with improvements in key health markers, but because of the need for recovery intervals after sprints, this potential can only truly be achieved if the number of sprint repetitions is low. Therefore, our observation that reducing the number of sprint repetitions will not attenuate the increase in $\dot{V}O_{2\max}$ associated with SIT, and in fact may possibly improve the effect, is an important novel finding.

Based predominantly on the results of studies investigating the dose–response relationship between regular aerobic exercise and improvements in health markers, it has generally been accepted that at a given exercise intensity, a greater volume of exercise training (in terms of training duration and frequency) is associated with greater improvements in $\dot{V}O_{2\max}$ (20). For example, in a clinical trial comparing low- or high-intensity aerobic training protocols with matched energy expenditure (Studies of a Targeted Risk Reduction Intervention through Defined Exercise I), the magnitude of change in $\dot{V}O_{2\max}$ was greater in the group exercising at a higher intensity (15). Although the volume of exercise used in HIIT and SIT protocols is generally reduced compared with aerobic exercise programs (11,47,63), the principle of a dose–response relationship has not been challenged in these studies directly; it is the interaction between training volume and intensity that is used to justify the lower volume. Thus, HIIT and SIT studies investigating the effects of protocols with a lower intensity or a shorter sprint duration tend to increase the number of sprint repetitions (45,69). Apart from two studies demonstrating that reducing sprint duration from 30 s to either 15 s (74) or 10 s (31) does not attenuate the improvement in $\dot{V}O_{2\max}$ with SIT, there have been no HIIT or SIT studies that have specifically investigated the dose–response relationship between volume of high-intensity exercise and the health outcomes. Our meta-analysis provides the first evidence that at “all-out” supramaximal exercise intensities the generally accepted positive association between volume of training and magnitude of training adaptations does not hold true. Thus, research into the health benefits of SIT should increase the focus on protocols with fewer sprints.

Because of the relatively low number of studies examining the effects of SIT protocols with fewer than six sprint repetitions, the present meta-analysis was not powerful enough to make conclusions on the optimal number of all-out sprint repetitions. Only two studies have investigated the effects of an SIT protocol incorporating just two sprints (50,52). As one of these used the largest sample size of all the studies included in the review ($n = 34$ [52]), the mean 10% increase in $\dot{V}O_{2\max}$ observed with this protocol (termed reduced-exertion high-intensity interval training [REHIT]) appears to be robust. The greatest improvement in absolute $\dot{V}O_{2\max}$ (17%) was reported by Gillen et al. (22), who modified the original REHIT protocol to include a third sprint. However, the total duration of this intervention was 12 wk, whereas at an intermediate measurement point after

TABLE 1. Training effects, training protocol parameters, and participant characteristics for the studies included in the meta-analysis.

Reference	Study Design	SIT Group Sample Size (n)	Proportion of Men	Mean Baseline $\dot{V}O_{2max}$ (ml·kg ⁻¹ ·min ⁻¹)	Mean Age (yr)	Mean BMI (kg·m ⁻²)	Training Duration (wk)	Total Training Sessions	Sprint Duration (s)	Recovery Duration (s)	Resistance (% of BM)	Sprint Repetitions		Effect on $\dot{V}O_{2max}$ (%)	
												Min	Max	Mean	SE
Metcalfe et al. (50)	C	11	0.45	34.2	25.0	23.5	6	18	20	200	7.5	1	2	12.7	2.8
Metcalfe et al. (52)	NC	34	0.50	35.0	34.1	24.6	6	18	20	200	5.0	1	2	9.6	1.5
Allemeier et al. (1)	C	11	1.00	48.7	22.7	24.8	6	15	30	1200	7.5	3	3	13.5	2.0
Gillen and Gibala (23)	NC	14	0.50	29.5	30.0	-	6	18	20	120	5.0	3	3	11.6	2.0
Ijichi et al. (35)	C	10	1.00	47.7	20.4	21.0	4	20	30	600	5.0	3	3	13.9	1.8
Gillen et al. (22)	C	9	1.00	32.0	27.0	27.0	12	36	30	120	5.0	3	3	17.3	3.3
Harris et al. (29)	C	6	0.00	35.0	22.0	23.6	4	12	20	270	7.5	4	4	9.0	3.4
Bayati et al. (8)	C	8	1.00	44.6	25.0	23.7	4	12	30	240	7.5	3	5	9.6	3.9
Barnett et al. (6)	C	8	1.00	47.6	20.4	-	8	24	30	180	-	3	6	8.2	2.1
Burgomaster et al. (11)	C	10	0.50	41.0	23.6	23.6	6	18	30	270	7.5	4	6	6.3	4.0
Hazell et al. (31)	C	13	0.81	47.0	24.0	24.7	2	6	30	240	10.0	4	6	8.3	2.2
Hazell et al. (31)	C	13	0.81	47.0	24.0	24.7	2	6	10	240	10.0	4	6	8.5	2.4
Hazell et al. (31)	C	13	0.81	47.0	24.0	24.7	2	6	10	120	10.0	4	6	3.9	1.3
Whyte et al. (76)	NC	10	1.00	32.8	32.1	30.3	2	6	30	270	6.5	4	6	8.4	2.6
Astorino et al. (2)	C	20	0.55	43.6	25.0	24.1	2	6	30	300	7.5	4	6	6.3	1.5
Shepperd et al. (63)	C	8	1.00	41.9	22.0	24.8	6	18	30	270	7.5	4	6	7.6	2.3
Larsen et al. (42)	NC	8	1.00	25.8	27.0	26.8	2	6	30	240	7.5	4	6	9.8	3.5
Ijichi et al. (35)	C	10	1.00	46.8	21.3	22.2	4	10	30	600	5.0	6	6	8.4	1.8
Kiviniemi et al. (40)	C	13	1.00	34.7	48.0	25.6	2	6	30	240	7.5	4	6	4.7	1.4
McGarr et al. (48)	C	8	0.75	47.2	25.0	-	2	8	30	240	7.5	4	6	14.2	4.5
Nalcaikan et al. (57)	C	8	1.00	40.2	21.7	25.5	7	21	30	270	7.5	4	6	7.0	1.8
Zelt et al. (77)	C	11	1.00	48.6	23.0	25.0	4	12	30	270	7.5	4	6	5.3	2.6
Zelt et al. (77)	C	12	1.00	43.9	22.0	26.0	4	12	15	285	7.5	4	6	7.4	2.7
Cochran et al. (13)	C	12	1.00	50.6	22.0	25.7	6	18	30	240	7.5	4	6	10.3	2.1
Burgomaster et al. (12)	C	8	0.75	44.6	22.0	25.6	2	6	30	240	7.5	4	7	1.4	2.0
Burgomaster et al. (10)	C	8	1.00	48.9	21.0	23.8	2	6	30	240	7.5	4	7	5.6	2.8
Bailey et al. (5)	C	8	0.63	42.0	21.0	23.7	2	6	30	240	7.5	3	7	7.4	2.4
Trilk et al. (70)	C	14	0.00	21.6	30.1	35.7	4	12	30	240	5.0	4	7	11.7	1.6
Richardson et al. (61)	C	9	0.56	40.0	21.0	23.8	2	6	30	240	7.5	4	7	11.2	2.7
Katz et al. (37)	NC	8	1.00	51.8	24.2	-	8	32	30	240	-	8	8	7.0	2.9
Scalzo et al. (62)	NC	21	0.52	41.5	22.5	22.4	3	9	30	240	7.5	4	8	3.7	2.1
Stathis et al. (68)	NC	8	0.75	49.6	22.1	-	7	21	30	180	-	3	10	4.2	2.9
McKenna et al. (49)	NC	8	1.00	47.1	20.9	23.7	7	21	30	180	7.5	4	10	13.7	3.2
MacDougall et al. (46)	NC	12	1.00	50.8	22.7	24.0	7	21	30	180	7.5	4	10	7.5	2.4
Harmer et al. (27)	NC	7	1.00	49.8	22.0	23.5	7	21	30	180	7.5	4	10	6.9	3.1
Harmer et al. (28)	C	7	0.71	43.7	24.0	23.8	7	21	30	180	7.5	4	10	8.2	3.6
Skleriyk et al. (64)	C	8	1.00	29.7	40.2	32.2	2	6	10	80	5.0	8	12	-1.7	2.9
Hellsten-Westling et al. (32)	NC	11	1.00	53.0	23.6	-	6	18	10	50	7.0	15	15	2.4	2.5

BM, body mass; BMI, body mass index; C, controlled; NC, not controlled.




FIGURE 3—Main effects of SIT on $\dot{V}O_{2max}$.

6 wk, the increase in $\dot{V}O_{2max}$ was 12%, very similar to the 10% and 14% improvements observed with the original REHIT protocol (50,52). Although future studies should determine whether the magnitude of the response for $\dot{V}O_{2max}$ is different between SIT protocols incorporating two to four sprints, the data presented in the present manuscript suggest that this difference will be small. If this is indeed the case, then several considerations support the use of the smallest number of sprints; that is, the two sprints used in the REHIT protocol. First, including a warm-up, recovery, and cooldown, this protocol has the potential to be the most time-efficient protocol. Furthermore, a drawback of

the use of SIT as a public health intervention is the potential for high associated perceived exertion and negative affective responses (8,21). In this light, it is important to point out that the number of sprint repetitions has been shown to negatively affect both of these parameters (19,44). Therefore, effective SIT protocols with fewer sprint repetitions will likely offer the best chance of sedentary target populations taking up and adhering to an SIT intervention for improving health (18). With this in mind, the available evidence suggests that two sprints can be recommended as effective at improving the important health marker of $\dot{V}O_{2max}$. It could be argued that considering the apparent linear association

TABLE 2. Main effect of SIT on $\dot{V}O_{2\max}$ and modifying effects.

	Effect on $\dot{V}O_{2\max}$ (Mean \pm 90% CL)	Inference
Main effect	7.8 \pm 4.0	Likely large increase
Modifying effects		
Two more sprint repetitions ^a	-1.2 \pm 0.8	Possibly small decrease
Three more training sessions ^a	0.7 \pm 0.4	Likely trivial change
10 s longer sprint duration ^a	0.6 \pm 1.3	Possibly trivial change
60 s longer recovery interval duration ^a	0.2 \pm 0.3	Most likely trivial change
3% of BM greater sprint resistance	1.0 \pm 2.3	Unclear
10 mL·kg ⁻¹ ·min ⁻¹ lower baseline $\dot{V}O_{2\max}$	1.5 \pm 1.9	Possibly small increase
7 yr higher age	-1.1 \pm 1.2	Possibly small decrease
6.2 kg·m ⁻² higher BMI	0.8 \pm 2.7	Unclear
Female sex	-0.2 \pm 3.5	Unclear
Uncontrolled study	-0.9 \pm 2.1	Unclear

The reference condition is an intervention using 14 SIT sessions and a maximum of seven repeated 30-s sprints with 240 s recovery. Effects of SIT are presented as the % change compared with pretraining. BMI, body mass index; CL, confidence limits; $\dot{V}O_{2\max}$, maximal aerobic capacity.

^aA practically relevant value was chosen to evaluate the effect magnitude; other numeric modifiers were evaluated as a 2 \times SD change in the parameter.

between the number of sprint repetitions and the improvement in $\dot{V}O_{2\max}$ (Fig. 4A), a single sprint could be expected to produce similar improvements with a lower time commitment. However, we have recently performed the first study to investigate the effects of a single supramaximal sprint on $\dot{V}O_{2\max}$ and observed no significant increase compared with a no-exercise control condition in response to 4 wk of training with a sample size of $n = 15$ (67). Further studies are required to confirm whether supramaximal sprints only improve $\dot{V}O_{2\max}$ if they are repeated. Furthermore, in light of the fact that the majority of studies that have studied the effects of SIT protocols incorporating two or three sprint repetitions have used 20-s sprints rather than the more commonly used 30-s sprints (22,23,50,52), further studies are required to investigate the shortest sprint duration that can be used without attenuating the adaptations to $\dot{V}O_{2\max}$.

Our present analysis does not provide an explanation for the possibly negative effect of reducing the maximal number of sprint repetitions on improvements in $\dot{V}O_{2\max}$, but a discussion of possible mechanisms is warranted. The main limiting factor of $\dot{V}O_{2\max}$ is generally assumed to be maximal cardiac output, possibly through increased blood volume (7,55). To date, no studies have examined the effect of SIT on blood

volume, but there is evidence in favor (17,74) and against (36) increases in blood volume in response to HIIT. Similarly, there is evidence in favor (3) and against (47) increased maximal cardiac output with SIT, with the latter finding suggesting that the adaptations to SIT for $\dot{V}O_{2\max}$ may be peripheral in origin. Indeed, several authors have proposed that improvements in $\dot{V}O_{2\max}$ with SIT are caused by improved skeletal muscle oxygen extraction because of increased mitochondrial density (22,36,57,65,77). Although it remains unclear whether the improvement in $\dot{V}O_{2\max}$ with SIT is due to central or peripheral adaptations, we propose that both increased blood volume and increased mitochondrial density could plausibly be explained by the rapid glycogen depletion associated with supramaximal exercise (51). First, maximal rates of glycogenolysis in the initial 15 s of a supramaximal sprint (58) are associated with the accumulation of metabolic derivatives, resulting in a hypertonic intramyocellular environment, influx of water, and a transient ~15%–20% drop in plasma volume within a time span of just a few minutes (51). This severe disturbance of circulatory homeostasis could be a stimulus for the body to increase blood volume in response to repeated SIT sessions. Second, glycogenolysis is associated with the release and activation of glycogen-bound 5' AMP-activated protein kinase (59), which through downstream signaling pathways involving peroxisome proliferator-activated receptor gamma coactivator 1- α (a proposed master regulator of aerobic adaptations) could be a mechanism leading to increased mitochondrial density (30). Glycogen breakdown during repeated supramaximal sprints has been shown to be completely attenuated by the time of the third sprint (58), and it is therefore plausible, for both of these speculated mechanisms, that performing just two repeated supramaximal sprints is sufficient to “saturate” (i.e., maximally activate) the adaptive response. In other words, if either increased blood volume or mitochondrial density underpins the changes in $\dot{V}O_{2\max}$ with SIT, and if rapid glycogen breakdown regulates those adaptations, then no additional improvements would be expected if more than 2–3 sprints are performed in a training session.

Apart from this hypothesis, it is also possible that increasing the number of sprint repetitions will result in “pacing” strategies that affect the “all-out” nature of the sprints (e.g.,




FIGURE 4—Modifying effects of number of sprint repetitions (A), baseline $\dot{V}O_{2\max}$ (B), and age (C) on the effect of SIT on $\dot{V}O_{2\max}$. Data points represent individual trials included in the meta-analysis, and the size of the data point is proportional to study weighting. Solid and dotted lines represent the effect of the moderator \pm 90% confidence limits, respectively.

reduction of peak and mean power output in initial sprints), or that accumulated fatigue may reduce the effectiveness of later sprints. Furthermore, the fact that increasing the number of sprint repetitions does not enhance the improvement in $\dot{V}O_{2\max}$ with SIT provides strong evidence against a role for the magnitude of the acute effects of supramaximal sprints on oxygen transfer, energy turnover, or total energy use, as part of the stimulus for improving $\dot{V}O_{2\max}$ with SIT, because for each of these factors the stimulus should be greater with more sprint repetitions.

Several limitations to the present meta-analysis should be noted. First, to be of use as a practical intervention for preventing and/or treating inactivity-related chronic disease, SIT interventions should also be effective at improving for example insulin sensitivity and glycemic control, blood pressure, blood lipid profile, and body composition. Therefore, one limitation is that only $\dot{V}O_{2\max}$ was included as an outcome measure in the present analysis. However, insufficient data for a meta-analysis are available for the effects of SIT on blood pressure (14,23,76), blood lipid profile (4,76), and body composition (69,76), but the effect of SIT on insulin sensitivity and glycemic control has received more attention (4,22,23,50,52,60,76). However, the methods used to assess the effects of SIT on these parameters have varied, with different studies using oral glucose tolerance tests (4,50,52,76), intravenous glucose tolerance tests (22), euglycemic hyperinsulinemic clamps (60), or continuous glucose monitoring (23). This means that a meta-analysis of the effects of the number of sprint repetitions in an SIT protocol on insulin sensitivity and glycemic control is also currently not feasible. Nonetheless, the improvements in insulin sensitivity and glycemic control observed to date with SIT protocols incorporating two (50) or three sprints (22,23) are encouraging.

Second, because of the number of available SIT studies, the power of our meta-analysis is insufficient to conclude with certainty that the modifying effect of the number of sprint repetitions is negative; i.e., it remains possible that in reality performing more sprints will result in the same improvements in $\dot{V}O_{2\max}$ (a chance of approximately 1 in 3). However, this is not of major importance to the significance of our findings: even “no effect” of the number of sprint repetitions would

lead to the logical conclusion that performing SIT protocols with more than 2 or 3 sprints is unnecessary for improving $\dot{V}O_{2\max}$ in sedentary individuals. On the basis of the present analysis, the chance that in reality the effect of performing more sprints is positive was calculated as 0.0%.

A final limitation of our meta-analysis is that only SIT interventions using all-out intensities were included. Optimizing time-efficient interventions aimed at improving general health requires consideration of various parameters, and exercise intensity is undoubtedly one of the key parameters affecting the effectiveness of HIIT and SIT protocols. However, because of the large range of intensities used in SIT and HIIT protocols (~80%–350% of $\dot{V}O_{2\max}$), we felt it was important to attempt to “control” for this variable in the present analysis by including only studies that used “all-out” cycling exercise. Nonetheless, there is a clear need for studies examining the effect of the number of sprint repetitions at lower exercise intensities; for example, in HIIT studies.

In conclusion, in the present meta-analysis, we demonstrate that SIT is possibly more effective at improving $\dot{V}O_{2\max}$ if fewer sprint repetitions are performed in a training session. Considering the proclaimed aim of SIT to provide a time-efficient alternative/adjunct to high-volume moderate-intensity aerobic exercise, this finding has important implications for the design of practical SIT interventions. We put forward that SIT research should move away from further characterizing the commonly used 4–7 × 30-s Wingate protocol and toward establishing acceptable and effective protocols. This will require more studies to examine the modifying effects of a range of training parameters (including number of sprint repetitions, sprint duration, sprint intensity, and training frequency) on adaptations to key health markers, as well as exercise enjoyment and acceptability, perceived exertion, and the potential to cause negative affective responses.

The authors thank Professor Will Hopkins for helpful advice during the preparation of the manuscript. No sources of funding were used to assist in the preparation of this review. The authors have no conflicts of interest to declare. The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. The results of the present study do not constitute endorsement by the American College of Sports Medicine.

REFERENCES

1. Allemeier CA, Fry AC, Johnson P, Hikida RS, Hagerman FC, Staron RS. Effects of sprint cycle training on human skeletal muscle. *J Appl Physiol* (1985). 1994;77(5):2385–90.
2. Astorino TA, Allen RP, Roberson DW, et al. Adaptations to high-intensity training are independent of gender. *Eur J Appl Physiol*. 2011;111(7):1279–86.
3. Astorino TA, Edmunds RM, Clark A, et al. High-intensity interval training increases cardiac output and $\dot{V}O_{2\max}$. *Med Sci Sports Exerc*. 2017;49(2):265–73.
4. Babraj JA, Vollaard NB, Keast C, Guppy FM, Cottrell G, Timmons JA. Extremely short duration high intensity interval training substantially improves insulin action in young healthy males. *BMC Endocr Disord*. 2009;9:3.
5. Bailey SJ, Wilkerson DP, Dimenna FJ, Jones AM. Influence of repeated sprint training on pulmonary O_2 uptake and muscle deoxygenation kinetics in humans. *J Appl Physiol* (1985). 2009;106(6):1875–87.
6. Barnett C, Carey M, Proietto J, Cerin E, Febbraio MA, Jenkins D. Muscle metabolism during sprint exercise in man: influence of sprint training. *J Sci Med Sport*. 2004;7(3):314–22.
7. Bassett DR Jr, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc*. 2000;32(1):70–84.
8. Bayati M, Farzad B, Gharakhanlou R, Agha-Alinejad H. A practical model of low-volume high-intensity interval training induces performance and metabolic adaptations that resemble ‘all-out’ sprint interval training. *J Sports Sci Med*. 2011;10(3):571–6.

9. Blair SN, Kohl HW 3rd, Barlow CE, Paffenbarger RS Jr, Gibbons LW, Macera CA. Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *JAMA*. 1995; 273(14):1093–8.
10. Burgomaster KA, Heigenhauser GJ, Gibala MJ. Effect of short-term sprint interval training on human skeletal muscle carbohydrate metabolism during exercise and time-trial performance. *J Appl Physiol (1985)*. 2006;100(6):2041–7.
11. Burgomaster KA, Howarth KR, Phillips SM, et al. Similar metabolic adaptations during exercise after low volume sprint interval and traditional endurance training in humans. *J Physiol*. 2008;586(1):151–60.
12. Burgomaster KA, Hughes SC, Heigenhauser GJ, Bradwell SN, Gibala MJ. Six sessions of sprint interval training increases muscle oxidative potential and cycle endurance capacity in humans. *J Appl Physiol (1985)*. 2005;98(6):1985–90.
13. Cochran AJ, Percival ME, Tricarico S, et al. Intermittent and continuous high-intensity exercise training induce similar acute but different chronic muscle adaptations. *Exp Physiol*. 2014; 99(5):782–91.
14. Cocks M, Shaw CS, Shepherd SO, et al. Sprint interval and endurance training are equally effective in increasing muscle microvascular density and eNOS content in sedentary males. *J Physiol*. 2013;591(3):641–56.
15. Duscha BD, Slentz CA, Johnson JL, et al. Effects of exercise training amount and intensity on peak oxygen consumption in middle-age men and women at risk for cardiovascular disease. *Chest*. 2005;128(4):2788–93.
16. Ekelund U, Ward HA, Norat T, et al. Physical activity and all-cause mortality across levels of overall and abdominal adiposity in European men and women: the European Prospective Investigation into Cancer and Nutrition Study (EPIC). *Am J Clin Nutr*. 2015; 101(3):613–21.
17. Esfandiari S, Sasson Z, Goodman JM. Short-term high-intensity interval and continuous moderate-intensity training improve maximal aerobic power and diastolic filling during exercise. *Eur J Appl Physiol*. 2014;114(2):331–43.
18. Foster C, Farland CV, Guidotti F, et al. The effects of high intensity interval training vs steady state training on aerobic and anaerobic capacity. *J Sports Sci Med*. 2015;14(4):747–55.
19. Frazão DT, de Farias Junior LF, Dantas TC, et al. Feeling of pleasure to high-intensity interval exercise is dependent of the number of work bouts and physical activity status. *PLoS One*. 2016;11(3):e0152752.
20. Garber CE, Blissmer B, Deschenes MR, et al. American College of Sports Medicine Position Stand: quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011; 43(7):1334–59.
21. Gillen JB, Gibala MJ. Is high-intensity interval training a time-efficient exercise strategy to improve health and fitness? *Appl Physiol Nutr Metab*. 2014;39(3):409–12.
22. Gillen JB, Martin BJ, MacInnis MJ, Skelly LE, Tarnopolsky MA, Gibala MJ. Twelve weeks of sprint interval training improves indices of cardiometabolic health similar to traditional endurance training despite a five-fold lower exercise volume and time commitment. *PLoS One*. 2016;11(4):e0154075.
23. Gillen JB, Percival ME, Skelly LE, et al. Three minutes of all-out intermittent exercise per week increases skeletal muscle oxidative capacity and improves cardiometabolic health. *PLoS One*. 2014; 9(11):e111489.
24. Gist NH, Fedewa MV, Dishman RK, Cureton KJ. Sprint interval training effects on aerobic capacity: a systematic review and meta-analysis. *Sports Med*. 2014;44(2):269–79.
25. Hallal PC, Andersen LB, Bull FC, et al. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet*. 2012;380(9838):247–57.
26. Hardcastle SJ, Ray H, Beale L, Hagger MS. Why sprint interval training is inappropriate for a largely sedentary population. *Front Psychol*. 2014;5:1505.
27. Harmer AR, McKenna MJ, Sutton JR, et al. Skeletal muscle metabolic and ionic adaptations during intense exercise following sprint training in humans. *J Appl Physiol (1985)*. 2000;89(5):1793–803.
28. Harmer AR, Ruell PA, McKenna MJ, et al. Effects of sprint training on extrarenal potassium regulation with intense exercise in Type 1 diabetes. *J Appl Physiol (1985)*. 2006;100(1):26–34.
29. Harris E, Rakobowchuk M, Birch KM. Sprint interval and sprint continuous training increases circulating CD34+ cells and cardiorespiratory fitness in young healthy women. *PLoS One*. 2014; 9(9):e108720.
30. Hawley JA. Molecular responses to strength and endurance training: are they incompatible? *Appl Physiol Nutr Metab*. 2009;34(3): 355–61.
31. Hazell TJ, Macpherson RE, Gravelle BM, Lemon PW. 10 or 30-s sprint interval training bouts enhance both aerobic and anaerobic performance. *Eur J Appl Physiol*. 2010;110(1):153–60.
32. Hellsten-Westing Y, Balsom PD, Norman B, Sjodin B. The effect of high-intensity training on purine metabolism in man. *Acta Physiol Scand*. 1993;149(4):405–12.
33. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc*. 2009;41(1):3–13.
34. Hunter DJ, Reddy KS. Noncommunicable diseases. *N Engl J Med*. 2013;369(14):1336–43.
35. Ijichi T, Hasegawa Y, Morishima T, Kurihara T, Hamaoka T, Goto K. Effect of sprint training: training once daily versus twice every second day. *Eur J Sport Sci*. 2015;15(2):143–50.
36. Jacobs RA, Fluck D, Bonne TC, et al. Improvements in exercise performance with high-intensity interval training coincide with an increase in skeletal muscle mitochondrial content and function. *J Appl Physiol (1985)*. 2013;115(6):785–93.
37. Katz A, Sharp RL, King DS, Costill DL, Fink WJ, et al. Effect of high intensity interval training on 2,3-diphosphoglycerate at rest and after maximal exercise. *Eur J Appl Physiol Occup Physiol*. 1984;52(3):331–5.
38. Keteyian SJ, Brawner CA, Savage PD, et al. Peak aerobic capacity predicts prognosis in patients with coronary heart disease. *Am Heart J*. 2008;156(2):292–300.
39. Kimm SY, Glynn NW, McMahon RP, Voorhees CC, Striegel-Moore RH, Daniels SR. Self-perceived barriers to activity participation among sedentary adolescent girls. *Med Sci Sports Exerc*. 2006;38(3):534–40.
40. Kiviniemi AM, Tulppo MP, Eskelinen JJ, et al. Autonomic function predicts fitness response to short-term high-intensity interval training. *Int J Sports Med*. 2015;36(11):915–21.
41. Korkiakangas EE, Alahuhta MA, Laitinen JH. Barriers to regular exercise among adults at high risk or diagnosed with type 2 diabetes: a systematic review. *Health Promot Int*. 2009;24(4):416–27.
42. Larsen RG, Befroy DE, Kent-Braun JA. High-intensity interval training increases in vivo oxidative capacity with no effect on P(i) → ATP rate in resting human muscle. *Am J Physiol Regul Integr Comp Physiol*. 2013;304(5):R333–42.
43. Lee DC, Sui X, Artero EG, et al. Long-term effects of changes in cardiorespiratory fitness and body mass index on all-cause and cardiovascular disease mortality in men: the Aerobics Center Longitudinal Study. *Circulation*. 2011;124(23):2483–90.
44. Little JP, Gillen JB, Percival ME, et al. Low-volume high-intensity interval training reduces hyperglycemia and increases muscle mitochondrial capacity in patients with type 2 diabetes. *J Appl Physiol (1985)*. 2011;111(6):1554–60.
45. Little JP, Safdar A, Wilkin GP, Tarnopolsky MA, Gibala MJ. A practical model of low-volume high-intensity interval training induces mitochondrial biogenesis in human skeletal muscle: potential mechanisms. *J Physiol*. 2010;588(Pt 6):1011–22.

46. MacDougall JD, Hicks AL, MacDonald JR, McKelvie RS, Green HJ, Smith KM. Muscle performance and enzymatic adaptations to sprint interval training. *J Appl Physiol (1985)*. 1998;84(6):2138–42.
47. Macpherson RE, Hazell TJ, Olver TD, Paterson DH, Lemon PW. Run sprint interval training improves aerobic performance but not maximal cardiac output. *Med Sci Sports Exerc*. 2011;43(1):115–22.
48. McGarr GW, Hartley GL, Cheung SS. Neither short-term sprint nor endurance training enhances thermal response to exercise in a hot environment. *J Occup Environ Hyg*. 2014;11(1):47–53.
49. McKenna MJ, Heigenhauser GJ, McKelvie RS, Obminski G, MacDougall JD, Jones NL. Enhanced pulmonary and active skeletal muscle gas exchange during intense exercise after sprint training in men. *J Physiol*. 1997;501(Pt 3):703–16.
50. Metcalfe RS, Babraj JA, Fawcner SG, Vollaard NB. Towards the minimal amount of exercise for improving metabolic health: beneficial effects of reduced-exertion high-intensity interval training. *Eur J Appl Physiol*. 2012;112(7):2767–75.
51. Metcalfe RS, Koumanov F, Ruffino JS, et al. Physiological and molecular responses to an acute bout of reduced-exertion high-intensity interval training (REHIT). *Eur J Appl Physiol*. 2015;115(11):2321–34.
52. Metcalfe RS, Tardif N, Thompson D, Vollaard NBJ. Health benefits of reduced-exertion high-intensity interval training (REHIT) are not different between sedentary men and women. *Appl Physiol Nutr Metab*. 2016;41(11):1117–23.
53. Milanovic Z, Sporis G, Weston M. Effectiveness of high-intensity interval training (HIT) and continuous endurance training for $\dot{V}O_{2\max}$ improvements: a systematic review and meta-analysis of controlled trials. *Sports Med*. 2015;45(10):1469–81.
54. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339:b2535.
55. Montero D, Diaz-Canestro C, Lundby C. Endurance training and $\dot{V}O_{2\max}$: role of maximal cardiac output and oxygen extraction. *Med Sci Sports Exerc*. 2015;47(10):2024–33.
56. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med*. 2002;346(11):793–801.
57. Nalcakan GR. The effects of sprint interval vs. continuous endurance training on physiological and metabolic adaptations in young healthy adults. *J Hum Kinet*. 2014;44:97–109.
58. Parolin ML, Chesley A, Matsos MP, Spriet LL, Jones NL, Heigenhauser GJ. Regulation of skeletal muscle glycogen phosphorylase and PDH during maximal intermittent exercise. *Am J Physiol*. 1999;277(5 Pt 1):E890–900.
59. Philp A, Hargreaves M, Baar K. More than a store: regulatory roles for glycogen in skeletal muscle adaptation to exercise. *Am J Physiol Endocrinol Metab*. 2012;302(11):E1343–51.
60. Richards JC, Johnson TK, Kuzma JN, et al. Short-term sprint interval training increases insulin sensitivity in healthy adults but does not affect the thermogenic response to beta-adrenergic stimulation. *J Physiol*. 2010;588(Pt 15):2961–72.
61. Richardson AJ, Gibson OR. Simulated hypoxia does not further improve aerobic capacity during sprint interval training. *J Sports Med Phys Fitness*. 2015;55(10):1099–106.
62. Scalzo RL, Peltonen GL, Binns SE, et al. Greater muscle protein synthesis and mitochondrial biogenesis in males compared with females during sprint interval training. *FASEB J*. 2014;28(6):2705–14.
63. Shepherd SO, Cocks M, Tipton KD, et al. Sprint interval and traditional endurance training increase net intramuscular triglyceride breakdown and expression of perilipin 2 and 5. *J Physiol*. 2013;591(3):657–75.
64. Skleryk JR, Karagounis LG, Hawley JA, Sharnan MJ, Laursen PB, Watson G. Two weeks of reduced-volume sprint interval or traditional exercise training does not improve metabolic functioning in sedentary obese men. *Diabetes Obes Metab*. 2013;15(12):1146–53.
65. Sloth M, Sloth D, Overgaard K, Dalgas U. Effects of sprint interval training on $\dot{V}O_{2\max}$ and aerobic exercise performance: a systematic review and meta-analysis. *Scand J Med Sci Sports*. 2013;23(6):e341–52.
66. Smith TB, Hopkins WG. Variability and predictability of finals times of elite rowers. *Med Sci Sports Exerc*. 2011;43(11):2155–60.
67. Songsorn P, Lambeth-Mansell A, Mair JL, et al. Exercise training comprising of single 20-s cycle sprints does not provide a sufficient stimulus for improving maximal aerobic capacity in sedentary individuals. *Eur J Appl Physiol*. 2016;16(8):1511–7.
68. Stathis CG, Febbraio MA, Carey MF, Snow RJ. Influence of sprint training on human skeletal muscle purine nucleotide metabolism. *J Appl Physiol (1985)*. 1994;76(4):1802–9.
69. Trapp EG, Chisholm DJ, Freund J, Boutcher SH. The effects of high-intensity intermittent exercise training on fat loss and fasting insulin levels of young women. *Int J Obes (Lond)*. 2008;32(4):684–91.
70. Trilk JL, Singhal A, Bigelman KA, Cureton KJ. Effect of sprint interval training on circulatory function during exercise in sedentary, overweight/obese women. *Eur J Appl Physiol*. 2011;111(8):1591–7.
71. Trost SG, Owen N, Bauman AE, Sallis JF, Brown W. Correlates of adults' participation in physical activity: review and update. *Med Sci Sports Exerc*. 2002;34(12):1996–2001.
72. Vandenberghe TJ, Hopkins WG. Effects of acute carbohydrate supplementation on endurance performance: a meta-analysis. *Sports Med*. 2011;41(9):773–92.
73. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Soft*. 2010;36(3):1–48.
74. Warburton DE, Haykowsky MJ, Quinney HA, et al. Blood volume expansion and cardiorespiratory function: effects of training modality. *Med Sci Sports Exerc*. 2004;36(6):991–1000.
75. Weston M, Taylor KL, Batterham AM, Hopkins WG. Effects of low-volume high-intensity interval training (HIT) on fitness in adults: a meta-analysis of controlled and non-controlled trials. *Sports Med*. 2014;44(7):1005–17.
76. Whyte LJ, Gill JM, Cathcart AJ. Effect of 2 weeks of sprint interval training on health-related outcomes in sedentary overweight/obese men. *Metabolism*. 2010;59(10):1421–8.
77. Zelt JG, Hankinson PB, Foster WS, et al. Reducing the volume of sprint interval training does not diminish maximal and submaximal performance gains in healthy men. *Eur J Appl Physiol*. 2014;114(11):2427–36.